



# Estradiol-Mediated Axogenesis of Hypothalamic Neurons Requires ERK1/2 and Ryanodine Receptors-Dependent Intracellular $\text{Ca}^{2+}$ Rise in Male Rats

Lucas E. Cabrera Zapata<sup>1</sup>, Mariana Bollo<sup>1</sup> and María Julia Cambiasso<sup>1,2\*</sup>

<sup>1</sup>Instituto de Investigación Médica Mercedes y Martín Ferreyra, INIMEC-CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina, <sup>2</sup>Cátedra de Biología Celular, Facultad de Odontología, Universidad Nacional de Córdoba, Córdoba, Argentina

## OPEN ACCESS

### Edited by:

Gabriela Alejandra Salvador,  
Universidad Nacional del Sur,  
Argentina

### Reviewed by:

Thomas Behnisch,  
Institutes of Brain Science, Fudan  
University, China  
Se-Young Choi,  
Seoul National University,  
South Korea

### \*Correspondence:

María Julia Cambiasso  
jcambiasso@immf.uncor.edu

**Received:** 15 January 2019

**Accepted:** 12 March 2019

**Published:** 02 April 2019

### Citation:

Cabrera Zapata LE, Bollo M and  
Cambiasso MJ  
(2019) Estradiol-Mediated  
Axogenesis of Hypothalamic Neurons  
Requires ERK1/2 and Ryanodine  
Receptors-Dependent Intracellular  
 $\text{Ca}^{2+}$  Rise in Male Rats.  
*Front. Cell. Neurosci.* 13:122.  
doi: 10.3389/fncel.2019.00122

17 $\beta$ -estradiol (E2) induces axonal growth through extracellular signal-regulated kinase 1 and 2 (ERK1/2)-MAPK cascade in hypothalamic neurons of male rat embryos *in vitro*, but the mechanism that initiates these events is poorly understood. This study reports the intracellular  $\text{Ca}^{2+}$  increase that participates in the activation of ERK1/2 and axogenesis induced by E2. Hypothalamic neuron cultures were established from 16-day-old male rat embryos and fed with astroglia-conditioned media for 48 h. E2-induced ERK phosphorylation was completely abolished by a ryanodine receptor (RyR) inhibitor (ryanodine) and partially attenuated by an L-type voltage-gated  $\text{Ca}^{2+}$  channel (L-VGCC) blocker (nifedipine), an inositol-1,4,5-trisphosphate receptor (IP<sub>3</sub>R) inhibitor (2-APB), and a phospholipase C (PLC) inhibitor (U-73122). We also conducted  $\text{Ca}^{2+}$  imaging recording using primary cultured neurons. The results show that E2 rapidly induces an increase in cytosolic  $\text{Ca}^{2+}$ , which often occurs in repetitive  $\text{Ca}^{2+}$  oscillations. This response was not observed in the absence of extracellular  $\text{Ca}^{2+}$  or with inhibitory ryanodine and was markedly reduced by nifedipine. E2-induced axonal growth was completely inhibited by ryanodine. In summary, the results suggest that  $\text{Ca}^{2+}$  mobilization from extracellular space as well as from the endoplasmic reticulum is necessary for E2-induced ERK1/2 activation and axogenesis. Understanding the mechanisms of brain estrogenic actions might contribute to develop novel estrogen-based therapies for neurodegenerative diseases.

**Keywords:** hypothalamic neurons, axogenesis, estradiol, ERK1/2,  $\text{Ca}^{2+}$  signaling, ryanodine receptors

## INTRODUCTION

For many years, estrogens have been recognized as one of the main orchestrators of the sexual differentiation of the brain, acting during critical periods of development to organize neural circuits in a way that determines the modulatory/activational effects of gonadal hormones in adulthood. Testosterone secreted by male rodent testes during development is aromatized in neurons to 17 $\beta$ -estradiol (E2), which displays multiple cellular processes that finally set the masculine phenotype (McCarthy, 2008; Wright et al., 2010). More recently, accumulating evidence

indicates that E2 is not only a reproductive hormone but also a brain-derived neuroprotective factor, coordinating multiple signaling mechanisms that protect the brain from neurodegenerative diseases, affective disorders and cognitive decline (Arevalo et al., 2015). These beneficial actions in the brain have positioned estrogens as promising therapeutic compounds against different brain pathologies such as Parkinson and Alzheimer diseases, schizophrenia, multiple sclerosis, stroke, neuroinflammation, among others (Dye et al., 2012; Villa et al., 2016; McGregor et al., 2017; Giatti et al., 2018).

Classical estrogenic actions are mediated *via* intracellular estrogen receptors (ERs) that function as ligand-activated transcription factors to regulate the expression of estrogen-responsive genes. Additionally, estrogens generate a wide diversity of rapid “non-classical” effects, which occur in a range from some seconds to a few minutes *via* membrane-initiated mechanisms (Arevalo et al., 2012), including the triggering of  $\text{Ca}^{2+}$  signals (Beyer and Raab, 1998; Picotto et al., 1999; Wong et al., 2012), and the activation of several signaling pathways, such as phospholipase C (PLC)/inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) and diacylglycerol (Le Mellay et al., 1997; Chaban et al., 2004), nitric oxide synthase/nitric oxide (Kelly and Levin, 2001), adenylate cyclase/AMPc/protein kinase A (PKA; Beyer and Karolczak, 2000), phosphoinositide-3 kinase (PI3K; Garcia-Segura et al., 2010), PKC, and extracellular signal-regulated kinase 1 and 2 (ERK1/2) cascades (Wu et al., 2005).

Currently, it is known that E2 prevents cell death, promotes neuronal survival, and enhances neuritogenesis and synaptic plasticity in the brain (Carroll and Pike, 2008; Spence et al., 2013; Acaz-Fonseca et al., 2014; Khan et al., 2015; Lai et al., 2017; Céspedes Rubio et al., 2018). The induction of neurite outgrowth by E2 was first demonstrated by Toran-Allerand (1976, 1980) and Toran-Allerand et al. (1983) working with organotypic explant cultures of the preoptic area, hypothalamus, and cerebral cortex. This neuritogenic effect of the hormone was then observed in other brain regions, both directly related and unrelated with reproduction (Nishizuka and Arai, 1981; Reisert et al., 1987; Cambiasso et al., 1995; Murphy and Segal, 1996). Hypothalamic neurons *in vitro* undergo several intermediate stages of development from unpolarized to fully polarized cells (Díaz et al., 1992). Most of the neuritogenic effects of E2 were demonstrated in polarized neurons (stage III of development), which are characterized by the presence of axon (Díaz et al., 1992; Cambiasso et al., 1995). Previous studies from our laboratory have shown that E2 induces axonal growth through ERK1/2 activation in hypothalamic neurons of male embryos *in vitro*. Both axogenesis (Cambiasso and Carrer, 2001) and ERK1/2 activation (Gorosito and Cambiasso, 2008) mediated by the hormone are dependent on a membrane-initiated mechanism since E2-bovine serum albumin (BSA; a membrane-impermeable conjugate of E2) was as effective as free E2. Interestingly, decreasing intracellular  $\text{Ca}^{2+}$  by the  $\text{Ca}^{2+}$ -chelator BAPTA-AM or blocking  $\text{Ca}^{2+}$ -dependent PKC isoforms by Ro 32-0432 significantly decreased these E2 effects (Gorosito and Cambiasso, 2008). These findings strongly suggested an important role for  $\text{Ca}^{2+}$  in E2-induced ERK1/2 pathway activation and axonal growth; however, the

results did not provide the mechanism of E2-induced  $\text{Ca}^{2+}$  signaling in hypothalamic neurons.

In this study, we found that E2 evoked activation of  $\text{Ca}^{2+}$  entry *via* L-type voltage-gated  $\text{Ca}^{2+}$  channels (L-VGCCs) and promoted  $\text{Ca}^{2+}$  release through ryanodine receptors (RyRs). This early  $\text{Ca}^{2+}$  response underlies E2-induced ERK1/2 activation and axogenesis in hypothalamic neurons. Altogether, these results bring new insights about the mechanism of brain estrogenic actions and might contribute to developing novel estrogen-based therapies for neurodegenerative diseases.

## MATERIALS AND METHODS

### Animals and Cell Cultures

Embryos were obtained from pregnant Wistar rats at embryonic day 16 (E16). The day of vaginal plug was set as E0. Experimental procedures for handling and sacrificing animals were approved by the Animal Care and Use Committee at our institution (CICUAL-IMMF, INIMEC-CONICET-UNC; Córdoba, Argentina) and followed the NIH guidelines for care and use of laboratory animals. The minimum number of animals required was used for these experiments and suffering was minimized. Primary neuronal and astroglial cultures were prepared as previously described in Cambiasso et al. (2000). Pregnant rats were sacrificed by cervical dislocation under  $\text{CO}_2$  anesthesia, and the fetuses were dissected from the uterus. The male fetuses used for cultures were identified by visualization of the spermatic artery on the developing testes. Ventromedial hypothalamic and mesencephalic regions were dissected out and stripped off the meninges for primary neuronal and glial cultures, respectively. At E16, the axogenic effect of E2 is contingent on the presence of astroglia (Cambiasso et al., 1995) or astroglia-conditioned media from a target region (Cambiasso et al., 2000; Cambiasso and Carrer, 2001; Brito et al., 2004). The basal medium (BM) was (1:1) DMEM:Ham's F12 Nutrient Mixture, supplemented with 0.043% l-alanyl-l-glutamine (GlutaMAX I), 0.15% glucose, 100 U/ml penicillin and 100  $\mu\text{g}/\text{ml}$  streptomycin. All cultures were raised under phenol red-free conditions to avoid “estrogen-like effects” (Berthois et al., 1986). For neuronal cultures, the dissociated cell suspension was seeded on different supports pre-coated with 1 mg/ml poly-D-lysine depending on the experiment: 60 mm  $\times$  15 mm dishes (Corning Life Science, Tewksbury, MA, USA) for protein assays, 25 mm coverslips (Assistent, Germany) for  $\text{Ca}^{2+}$  imaging, and 12 mm coverslips (Assistent, Germany) for morphological studies.

### Western Blot

Hypothalamic neurons derived from male fetuses plated 1–2 h before were fed with astroglia-conditioned media for 2 days *in vitro* (DIV). After a 2 h washout period using BM, neuronal cultures were treated for 1 h with nifedipine (2  $\mu\text{M}$ ; Sigma-Aldrich, St. Louis, MO, USA), inhibitory ryanodine (50  $\mu\text{M}$ ; Santa Cruz Biotechnology, Santa Cruz, CA, USA), 2-APB (100  $\mu\text{M}$ ; Santa Cruz Biotechnology, Santa Cruz, CA, USA) or U-73122 (10  $\mu\text{M}$ ; Sigma-Aldrich, St. Louis, MO, USA), and then pulsed with 100 nM E2 (Sigma-Aldrich, St. Louis, MO, USA)

for 15 min. Hormone concentration used was determined by dose dependence (1–100 nM) experiments previously performed by our group (Gorosito and Cambiasso, 2008; Gorosito et al., 2008). ERK phosphorylation was maximally increased after the application of 100 nM E2. This dose was then used for all further acute stimulation studies. We have used compounds at final concentrations that did not alter cell viability or morphology in control conditions.

After treatment, hypothalamic neurons were washed and harvested at 4°C in RIPA buffer [150 mM NaCl, 0.1% NP40, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate (SDS), 50 mM Tris, pH 7.5] with protease and phosphatase inhibitors (1 µg/ml aprotinin, 1 µg/ml leupeptin, 1 µg/ml pepstatin A, 5 µg/ml chymostatin, 5 µg/ml antipain, 100 µg/ml PMSF, 50 µM NaF, 10 µM  $\text{Na}_4\text{P}_2\text{O}_7$ , and 1 mM  $\text{NaVO}_4$ ). Protein samples (20 µg/lane) were separated by 10% SDS-PAGE and transferred onto polyvinylidene fluoride membrane (Bio-Rad, Hercules, CA, USA). ERK1/2 phosphorylation was detected by using a rabbit monoclonal anti-phospho-p44/42 MAPK (Cell Signaling Technology, Danvers, MA, USA), which specifically detects both Thr202 and Tyr204 ERK phosphorylation forms of ERK [molecular masses (kDa) for ERK1 and ERK2 are 44 and 42, respectively]. Total ERK1/2 was detected by using a mouse anti-p44/42 MAPK (Cell Signaling Technology, Danvers, MA, USA). Secondary antibodies conjugated to horseradish peroxidase (Jackson, West Grove, PA, USA) were used for the detection by enhanced chemiluminescence on X-ray film. After incubation with the antibody against phospho-ERK1/2, blots were stripped and then re-probed with anti-total ERK1/2 to ensure equal protein loading. The resulting film samples were scanned and analyzed with an image analysis program (ImageJ; NIH, Bethesda, MD, USA). Data are presented as a ratio of phospho-ERK1/2/total-ERK1/2 of 3–4 different experiments (independent cultures) performed in duplicate.

## Ca<sup>2+</sup> Imaging

After 2 DIV, hypothalamic neurons were treated (or not) with 50 µM ryanodine or 2 µM nifedipine for 1 h and incubated with 3 µM acetoxymethyl (AM) ester form of the organic  $\text{Ca}^{2+}$ -dye Cal-520 (AAT Bioquest, Sunnyvale, CA, USA) for 30 min at 37°C in a  $\text{Ca}^{2+}$ -containing HEPES buffered salt solution ( $\text{Ca}^{2+}$ -HBSS) composed of (mM): 135 NaCl, 5.4 KCl, 2  $\text{CaCl}_2$ , 1  $\text{MgCl}_2$ , 10 HEPES, and 10 glucose; pH = 7.4 set with NaOH at RT. Following loading, neurons were washed twice with warm  $\text{Ca}^{2+}$ -HBSS and were imaged in the same buffer. To analyze the participation of extracellular  $\text{Ca}^{2+}$ , the cells were resuspended in an EGTA-containing buffer composed of (mM): 135 NaCl, 5 KCl, 0.5  $\text{CaCl}_2$ , 1.2  $\text{MgCl}_2$ , 5 HEPES, 14  $\text{NaHCO}_3$ , and 1 EGTA; pH = 7.4 set with NaOH at RT.

Imaging of cytosolic  $\text{Ca}^{2+}$  signals was performed using a 60× oil immersion objective of an Olympus IX81 inverted microscope [equipped with a Disk Spinning Unit (DSU), epifluorescence illumination (150 W Xenon Lamp), and a microprocessor], an ORCA AG (Hamamatsu) CCD camera and OSIS software. Frames were collected at a continuous rate of 2.5 per second during 5 min (790 frames). Cal-520 was excited at a wavelength of 492 nm, and emitted fluorescence was collected at 514 nm.

E2 (100 nM) was added 30 s after starting the recording. 10 µM thapsigargin (tg) was added at 3 min of recording as a positive control of normal endoplasmic reticulum  $\text{Ca}^{2+}$  content. The fluorescence intensity of the  $\text{Ca}^{2+}$  indicator was analyzed using ImageJ (NIH, Bethesda, MD, USA) software and plotted as the change in fluorescence ( $\Delta F$ ) of  $2 \times 2$  pixels divided by mean resting fluorescence [ $(F_0; \Delta F/F_0)$ ] over time. We measured both the peak fluorescence value and the integrated area under the  $\Delta F/F_0$  curve with OriginPro 8 SR0 software (OriginLab Corporation, Northampton, MA, USA). The integrated area roughly corresponds to the total amount of  $\text{Ca}^{2+}$  released over the recording period.


## Immunocytochemical Staining

To analyze the effect of ryanodine in E2-stimulated axon growth without affecting the normal polarization of neurons, we performed the experiment after 1 DIV (stage III of development). After 2 h in absence of E2, the cultures were treated for 1 h with 50 µM ryanodine before the addition of 10 nM E2 for an additional 24 h. The hormone concentration used to study the neuritogenic effect of E2 was chosen based on previous studies of our laboratory (Gorosito et al., 2008; Scerbo et al., 2014).

After 2 DIV, neuronal cultures were fixed for 20 min with warm 4% paraformaldehyde in PBS containing 0.12 M sucrose and rinsed in PBS. Neurons were immunocytochemically stained with antibodies against  $\beta$ -tubulin class III (SDL3D10). The details of the immunocytochemical procedure were as specified by Díaz et al. (1992). Briefly, the fixed cells were permeabilized in 0.2% Triton X-100 for 5 min at RT, preincubated with 5% BSA, incubated in mouse anti- $\beta$ -tubulin class III (Sigma-Aldrich, St. Louis, MO, USA), rinsed in PBS, and finally incubated with appropriate biotinylated secondary antibody. Incubation with secondary antibody was followed by washing in PBS, incubation for 2 h in VECTASTAIN ABC immunoperoxidase reagent (Vector Laboratories, Burlingame, CA, USA), and a final reaction with 1.4 mM 3,3'-diaminobenzidine in phosphate buffer with  $\text{H}_2\text{O}_2$ . Coverslips were then dehydrated with ethanol, cleared with xylene, and mounted on glass slides for morphometric analysis. No immunostaining was detected when primary antibodies were replaced by 5% BSA.

## Morphometric Analysis

The morphometric analysis of stained neuronal cultures was performed on digitized video images using JAVA as an image processor (Jandel Inc., Richmond, CA, USA) controlled by a host computer. Images were acquired through an optic microscope (Carl Zeiss, Germany). Microscope slides were coded, and the person conducting the analysis was blind to the experimental group. All labeled cells that could be identified as one individual neuron were measured in random fields at 40× magnification. Neural processes were classified as minor processes or axons according to accepted morphological criteria (Dotti et al., 1988; Blanco et al., 1990; Díaz et al., 1992). Minor processes are two or three short neurites that emerge from the cell body; axons are much longer, unique, thin, and relatively uniform in diameter. Neurons were considered to have developed an axon



**FIGURE 1 |** E2-induced extracellular signal-regulated kinase 1 and 2 (ERK1/2) activation depends on cytosolic  $\text{Ca}^{2+}$  increase mainly mediated by ryanodine receptors (RyRs). Effects of (A) 2  $\mu\text{M}$  nifedipine, (B) 50  $\mu\text{M}$  ryanodine, (C) 100  $\mu\text{M}$  2-APB or (D) 10  $\mu\text{M}$  U-73122 on E2-induced ERK phosphorylation. After washing for 2 h, the cultures were treated with the inhibitors for 1 h and were then pulsed for 15 min with 17 $\beta$ -estradiol (E2) and harvested for Western blotting. Top: ratio of readings for pERK/ERK bands in arbitrary densitometric units. Bottom: examples of immunoblots showing a decrease of hormone-induced ERK phosphorylation in cultures pretreated with the inhibitors. Molecular masses (kDa) for ERK1 and ERK2 are 44 and 42, respectively. Blots shown are representative of the mean  $\pm$  SEM of 3–4 different cultures. (A) Nifedipine: ANOVA  $F_{(2,10)} = 52.78$ ;  $p \leq 0.001$ . Least significant differences (LSDs) test indicated \*\*\* $p < 0.001$  vs. control and ♦♦ $p < 0.01$  vs. E2. (B) Ryanodine: ANOVA  $F_{(2,6)} = 5.856$ ;  $p = 0.04$ . LSDs test indicated \* $p = 0.05$  vs. control and ♦ $p = 0.05$  vs. E2. (C) 2-APB: ANOVA  $F_{(2,6)} = 27.203$ ;  $p \leq 0.001$ . LSDs test indicated \*\*\* $p < 0.001$  and \* $p = 0.05$  vs. control and ♦♦ $p = 0.01$  vs. E2. (D) U-73122: ANOVA  $F_{(2,6)} = 34.891$ ;  $p \leq 0.001$ . LSDs test indicated \*\*\* $p < 0.001$  and \*\* $p = 0.01$  vs. control and ♦ $p \leq 0.05$  vs. E2.

if they showed one neurite three to five times longer than the rest (stage III of development). Soma area, length of minor processes, total axonal length, and the number of neurites per cell were recorded. At least 60 neurons were measured for every experimental condition in each culture; at least three separate cultures were made for every condition.

## Statistical Analysis

Data were statistically evaluated by one-way ANOVA, followed by Fisher's Least Significant Difference (LSD) *post hoc* test


(Statistica; StatSoft Inc., Tulsa, OK, USA) where  $p < 0.05$  was considered statistically significant.

## RESULTS

### E2-Induced ERK1/2 Activation Is Mainly Mediated by RyRs

Our previous results suggested that E2-induced ERK1/2 phosphorylation is  $\text{Ca}^{2+}$ -dependent. Here, we further






**FIGURE 2 |** 17 $\beta$ -estradiol (E2) induces  $\text{Ca}^{2+}$  oscillations. **(A)** Hypothalamic neurons were loaded with Cal-520 AM for 30 min at 37°C, maintained in a  $\text{Ca}^{2+}$ -containing buffer ( $\text{Ca}^{2+}$ -HBSS) and changes in cytosolic  $\text{Ca}^{2+}$  concentration were measured using confocal microscopy [Olympus IX81 inverted microscope equipped with a Disk Spinning Unit (DSU)]. Time series of Cal-520 pseudocolor images is shown, before and after the addition of E2 and thapsigargin (tg). Pseudocolor scale bar: 0.3–0.001 arbitrary units. Length scale bar: 20  $\mu\text{m}$ . **(B)** Representative  $\text{Ca}^{2+}$  traces [regions of interest labeled 1–4 in **(A)**] plotted as changes over time in fluorescence intensity of the indicator ( $\Delta F$ ) respect to resting values ( $F_0$ ). Arrows indicate the addition time of E2 (30 s) and tg (3 min). Data are from one representative experiment out of six independent experiments.

investigated the  $\text{Ca}^{2+}$  response involved in ERK activation mediated by E2. Hypothalamic cultures grown with E2 for 48 h were washed in BM for 2 h and pre-treated with specific compounds for 1 h before a pulse of E2 for 15 min. In agreement with previous reports, E2 induced a strong phosphorylation of ERK at 15 min (**Figure 1**). This effect was completely abolished by inhibitory ryanodine (**Figure 1B**) and partially attenuated by nifedipine (**Figure 1A**),  $\text{IP}_3\text{R}$  inhibitor 2-APB (**Figure 1C**), and a PLC inhibitor U-73122 (**Figure 1D**).

### E2 Induces Rapid $\text{Ca}^{2+}$ Increase Depending on $\text{Ca}^{2+}$ Influx and RyRs

As E2-activation of the ERK1/2 signaling cascade depends on extracellular as well as intracellular  $\text{Ca}^{2+}$  stores, we decided to characterize the  $\text{Ca}^{2+}$  signal generated by the hormone. The addition of agonist to cell cultures, loaded with the indicator Cal-520 AM and imaged in a  $\text{Ca}^{2+}$ -containing buffer, induced fluorescence changes that were observed in neuronal soma as well as in minor processes (**Figure 2**). Seventeen out of



**FIGURE 3 |** Ryanodine, external  $\text{Ca}^{2+}$  and nifedipine modulate E2-induced  $\text{Ca}^{2+}$  increase. **(A)** Mean values of maximal  $\Delta F/Fo$  for  $\text{Ca}^{2+}$  mobilized by 17 $\beta$ -estradiol (E2) in hypothalamic neurons recorded either in the presence or absence of extracellular  $\text{Ca}^{2+}$  ( $\text{Ca}^{2+}$ -HBSS/EGTA-containing buffer, named High  $\text{Ca}^{2+}$ /Low  $\text{Ca}^{2+}$ ), or after a pre-incubation period (1 h) with ryanodine or nifedipine. ANOVA  $F_{(3,41)} = 22.94$ ;  $p \leq 0.001$ . LSDs test indicated \*\*\* $p < 0.001$  vs.  $\text{Ca}^{2+}$ -HBSS. **(B)** Mean values of integrated area under  $\Delta F/Fo$  curve (AUC) after thapsigargin (tg) addition during  $\text{Ca}^{2+}$  imaging [same conditions as **(A)**]. ANOVA  $F_{(3,26)} = 9.18$ ;  $p \leq 0.001$ . LSDs test indicated \*\*\* $p < 0.001$  and \*\* $p = 0.01$  vs. ryanodine. Bars represent mean  $\pm$  SEM;  $n = 4$ –6 different cultures.

76 neurons (22.4%) from six independent experiments imaged in  $\text{Ca}^{2+}$ -HBSS responded to E2 pulses within 15–100 s of treatment (average =  $46.8 \pm 6.60$  s). These  $\text{Ca}^{2+}$  events often occur in repetitive oscillations, which display interspike intervals of  $21.58 \pm 8.83$  s (9 out of 17 neurons). The amplitude in terms of  $\Delta F/Fo$  was  $0.154 \pm 0.01$ . The total amount of  $\text{Ca}^{2+}$  mobilized, measured as the integrated area under the curve (AUC) for all E2-generated peaks, was  $4.69 \pm 1.03$ , which was 90.6% with respect to the control [remained endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA) inhibitor].

This E2-induced  $\text{Ca}^{2+}$  signal was not observed in the absence of extracellular  $\text{Ca}^{2+}$ . Moreover, we found that pre-incubation of neuronal cultures with inhibitory ryanodine suppressed E2-evoked  $\text{Ca}^{2+}$  release (Figure 3A). Importantly, under these conditions, the tg was able to mobilize amounts of  $\text{Ca}^{2+}$ , measured as AUC, of  $14.69 \pm 3.45$  (EGTA-containing buffer) and  $35.05 \pm 11.54$  (ryanodine pre-incubation), five and nine times greater, respectively, than the amounts of  $\text{Ca}^{2+}$  mobilized by tg in E2-induced control neurons (Figure 3B). Moreover, nifedipine reduced the E2-induced  $\text{Ca}^{2+}$  increase more than 50% ( $\Delta F/Fo = 0.065 \pm 0.011$ ; and  $\text{AUC} = 10.779 \pm 1.159$ ,  $n = 4$ ), which strongly suggests the participation of L-VGCCs in this signal (Figures 3A,B). Representative  $\text{Ca}^{2+}$  traces plotted as  $\Delta F/Fo$  vs. time for EGTA, ryanodine, and nifedipine conditions are provided in Supplementary Figure S1.

Taken together, these results indicate that both  $\text{Ca}^{2+}$  influx and mobilization from intracellular stores contribute to the response.

## E2-Induced Axonal Growth Depends on $\text{Ca}^{2+}$ Signal Generated by RyRs


Finally, we tested whether the RyR-induced  $\text{Ca}^{2+}$  response is part of the signaling cascade that mediates the axogenic effect

of E2. Neurons grew under the following conditions: in the presence and absence of E2 and pretreated with inhibitory ryanodine. After these treatments, the cells were grown for an additional 24 h period with (E2) or without E2 (control). In agreement with previous results, the morphometric analysis indicated that neurons grown under hormonal treatment show longer axons than neurons in control conditions without E2 (Figure 4A). Remarkably, blocking RyRs with ryanodine completely inhibited the E2-induced axogenesis (Figure 4B). Moreover, no significant differences were observed in the number of primary neurites, length of minor processes, or soma area per neuron resulting from E2 or ryanodine treatment (Table 1), confirming that the hormonal effect is restricted to axonal growth (Díaz et al., 1992; Cambiasso et al., 1995, 2000; Brito et al., 2004).

## DISCUSSION

In the present study, results from ERK1/2 phosphorylation,  $\text{Ca}^{2+}$  imaging and neuronal growth consistently pointed to RyRs as the  $\text{Ca}^{2+}$  channels necessary to mediate activation of the MAPK/ERK pathway and the final axogenic effect induced by E2. Inhibitory ryanodine completely blocked E2-mediated  $\text{Ca}^{2+}$  transients, ERK1/2 phosphorylation and axonal outgrowth, which provides valuable evidence to propose that E2 mobilizes endoplasmic reticulum stores of  $\text{Ca}^{2+}$  through RyRs to activate the signaling cascades that finally affect the axonal elongation of hypothalamic neurons.

In regard to the impact of functional RyRs on cellular response, Dr Hidalgo's group showed that BDNF-induced neural plasticity requires functional RyRs activated by the  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR) mechanism to evoke the larger  $\text{Ca}^{2+}$  signaling needed to maintain changes during long-term memory storage (Adasme et al., 2011). CICR is a positive



**FIGURE 4 |** RyR activity is required to the E2-induced axogenesis. **(A)** Representative images of male hypothalamic neurons cultured for 48 h with (E2) or without (control) 10 nM 17 $\beta$ -estradiol in combination or not with 50  $\mu\text{M}$  ryanodine (Ry) for the last 24 h of incubation (arrows indicate the axons of some neurons). **(B)** Mean of axonal length for each condition in **(A)**. ANOVA  $F_{(3,12)} = 4.51$ ;  $p = 0.02$ . LSDs test indicated \* $p = 0.05$  vs. control and ♦♦ $p = 0.01$  vs. E2. Data represent the mean  $\pm$  SEM;  $n = 4$  independent cultures. Scale bar: 100  $\mu\text{m}$ .

**TABLE 1 |** Number of primary neurites, soma area, and length of minor processes of male hypothalamic neurons grown with or without 17 $\beta$ -estradiol (E2) in 3 combination or not with ryanodine.

Variable	Treatment			
	Control	E2	E2 + Ryanodine	Ryanodine
N° of neurites	4.2 $\pm$ 0.2	4.4 $\pm$ 0.2	4.3 $\pm$ 0.4	4.2 $\pm$ 0.2
Soma area ( $\mu\text{m}^2$ )	118.9 $\pm$ 13.2	116.6 $\pm$ 4.5	119.5 $\pm$ 10.0	126.2 $\pm$ 9.0
Minor processes length ( $\mu\text{m}$ )	107.2 $\pm$ 7.7	115.6 $\pm$ 8.9	94.0 $\pm$ 11.5	92.6 $\pm$ 5.8

Data represent the mean  $\pm$  SEM;  $n = 4$  independent cultures.

feedback mechanism by which cytoplasmic  $\text{Ca}^{2+}$  stimulates  $\text{Ca}^{2+}$  release from the endoplasmic reticulum through RyRs or IP<sub>3</sub>R (Bezprozvanny et al., 1991; Berridge et al., 2003; Seo et al., 2015).

Moreover, several studies have indicated a regulatory role for estrogens on RyRs activity, for instance, in the human eccrine sweat gland cell line NCL-SG3 (Muchekehu and Harvey, 2008),

ventricular myocytes (Yan et al., 2011), detrusor smooth muscle cells (Hristov et al., 2017), and dorsal root ganglion neurons (Ferrari et al., 2016; Khomula et al., 2017). Interestingly, Zhao X. et al. (2005) proposed, in a neuroblastoma cell line, an E2-mediated mechanism starting at the plasma membrane, by which rapid  $\text{Ca}^{2+}$  signaling potentiates the transcription of genes normally regulated by estrogens; RyRs,  $\text{IP}_3$ Rs, and N-VGCCs, but not L-VGCCs, were involved in the process. Our results clearly show that RyRs are indispensable  $\text{Ca}^{2+}$  channels involved in the non-classical signaling events produced by E2 to generate axonal growth in hypothalamic neurons. However, RyRs also require a previous small increase in cytosolic  $\text{Ca}^{2+}$  from resting levels to activate and release  $\text{Ca}^{2+}$  by CICR (Hidalgo et al., 2005; Lanner et al., 2010). Since we found that removing extracellular  $\text{Ca}^{2+}$  or blocking the membrane channels L-VGCCs prevented the  $\text{Ca}^{2+}$  signaling induced by E2 and that inhibiting L-VGCCs also reduced ERK1/2 phosphorylation modulated by the steroid, we postulate that E2 initially induces a  $\text{Ca}^{2+}$  influx in hypothalamic neurons via L-VGCCs that then enables RyRs opening to generate the final and complete  $\text{Ca}^{2+}$  signaling event (Hidalgo, 2005; Calin-Jageman and Lee, 2008). Besides L-VGCCs, our results show that the PLC/ $\text{IP}_3$ Rs system is involved in E2-induced ERK1/2 activation, since both 2-APB and U-73122 used as blockers of  $\text{IP}_3$ Rs and PLC, respectively, produced a significant reduction in phosphorylation levels of the kinases in the presence of the hormone. The activation of PLC and  $\text{Ca}^{2+}$  release via  $\text{IP}_3$ Rs induced by estrogens has been previously reported in different cellular systems (Chaban et al., 2004; Fricke et al., 2007).

L-VGCCs are the major route of  $\text{Ca}^{2+}$  entry into neurons and the most profusely studied and best characterized VGCC type by far, as they play a predominant role in the brain (Striessnig et al., 2014; Vega-Vela et al., 2017). Several studies report that E2 is able to modulate L-VGCCs activity (Bulayeva et al., 2005; Sarkar et al., 2008; Farkas et al., 2012; Feng et al., 2013). Wu et al. (2005) and Zhao L. et al. (2005) indicated that E2 induced rapid  $\text{Ca}^{2+}$  influx through L-VGCCs, which was required to activate the Src/ERK/CREB/Bcl-2 signaling pathway and finally mediated neuroprotective and neurotrophic responses in rat hippocampal and cortical neurons. The generation of this intracellular  $\text{Ca}^{2+}$  increase and the downstream activation of ERK depend on the presence of ERs in the membrane of rat hippocampal neurons (Wu et al., 2011). These membrane ER-expressing neurons represented 29% of the cultured cells and all of them co-expressed L-VGCCs. Consistently, our data from  $\text{Ca}^{2+}$  imaging experiments indicated that approximately 22% of hypothalamic neurons responded to E2.

Although previously it has been reported that E2 induces L-VGCCs-mediated  $\text{Ca}^{2+}$  influx, the question about how the hormone activates L-VGCCs remains open. Sarkar et al. (2008) reported that E2 potentiated the activity of L-VGCCs by directly binding to specific sites in the channel, independently of ERs. On the other hand, PI3K signaling cascade has been proposed as a candidate to link membrane ER activation with L-VGCCs aperture (Simoncini et al., 2000; Quignard

et al., 2001; Wu et al., 2005), although our preceding work blocking PI3K with LY-294,002 did not prevent the axogenic effect of E2 in hypothalamic neurons (Gorosito and Cambiasso, 2008). It is important to note that we have previously reported the expression of ER $\alpha$  on the cell-surface of embryonic hypothalamic neurons (Gorosito et al., 2008), and that the membrane-impermeable E2-albumin construct (E2-BSA) was as effective as free E2 to generate ERK1/2 phosphorylation (Gorosito and Cambiasso, 2008) and axonal elongation (Cambiasso and Carrer, 2001), evidence that altogether indicate these processes respond to a membrane-initiated ER $\alpha$ -mediated mechanism.

Another question that arises is what other elements lead from the RyRs-mediated  $\text{Ca}^{2+}$  release to ERK1/2 activation. In Gorosito and Cambiasso (2008) we reported that PKC but not PKA nor CaMKII is required in the E2-induced MAPK-ERK pathway activation, since an inhibitor with specificity for the PKC  $\text{Ca}^{2+}$ -dependent  $\alpha$  and  $\beta$ I isoforms, Ro 32-0432, attenuated E2-modulated ERK1/2 phosphorylation and prevented the axogenic effect of the hormone. PKC activation by E2 has been found in different cell types, including breast cancer cells, hepatocytes, and cortical and hypothalamic neurons (Marino et al., 1998; Boyan et al., 2003; Cordey et al., 2003; Qiu et al., 2003).  $\text{Ca}^{2+}$ -dependent PKC activation can then induce ERK1/2 phosphorylation via Src/Ras signaling (Cullen and Lockyer, 2002; Brandt et al., 2003; Roskoski, 2005).

Our results show that E2-induced cytosolic  $\text{Ca}^{2+}$  often increases as repetitive oscillations. This is in agreement with previous work demonstrating E2-induced intracellular  $\text{Ca}^{2+}$  oscillations that involved internal stores and PKA and PLC activity in neurons of the arcuate nucleus (Fricke et al., 2007). It is well known that the frequency of  $\text{Ca}^{2+}$  oscillation may depend on  $\text{Ca}^{2+}$  influx into the cell (Sneyd et al., 2004), SERCA activity (Falcke et al., 2003) and oscillating cytoplasmic  $\text{IP}_3$  concentration (Sneyd et al., 2006). Our data suggest that the frequency of  $\text{Ca}^{2+}$  oscillations in hypothalamic neurons stimulated by E2 mainly depends on  $\text{Ca}^{2+}$  entry. Leaving aside its modulation, our results allow us to speculate that the characteristic frequency encodes information to regulate the cellular response (axonal growth) mediated by the hormone (Dolmetsch et al., 1998).

In conclusion, we have provided new insights into the non-classical mechanisms triggered by estrogens and its axogenic effect in male rat hypothalamic neurons. The hormone induces ERK1/2 activation in a  $\text{Ca}^{2+}$ -dependent manner. RyRs inhibition abolished this activation as well as axonal growth. The oscillatory  $\text{Ca}^{2+}$  signal generated by E2 required functional RyRs and L-VGCCs. This early  $\text{Ca}^{2+}$  response that underlies E2-induced RyRs and MAPK/ERK activation may transmit a finely tuned message into a neuronal development program, reflecting the need for tight control of a critical event during sexual differentiation of the male brain. The conjunction of XY genotype with adequate estrogen exposure levels at the time of hypothalamic neuronal differentiation may induce the growth of axons towards their appropriate targets. A complete and detailed understanding of the intracellular signaling mechanisms and neuronal processes mediated by estrogens will allow to



improve current estrogen-based therapies, such as hormone replacement therapy in postmenopausal women, as well as to develop novel treatments to prevent and/or alleviate neurological pathologies based on its widely proven neuritogenic and neuroprotective effects.

## AUTHOR CONTRIBUTIONS

LCZ, MB and MC made the conception and design of research, interpreted results of experiments, wrote the manuscript. LCZ performed all experiments and analyzed the data. LCZ and MB prepared the figures. MB and MC edited and revised the manuscript. MC drafted the manuscript.

## FUNDING

This study was supported by grants from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET;

PIP 2013-2015), Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT; PICT 2015 N° 1333) and Secretaría de Ciencia y Tecnología (SECyT)-Universidad Nacional de Córdoba (2016-2017), Argentina to MC and from CONICET (PIP 2014-2016) to MB.

## ACKNOWLEDGMENTS

We thank for the technical support from INIMEC animal facility and Dr Quassollo for his invaluable help with imaging. Calcium imaging was generated at National Center of Microscopy, Universidad Nacional de Córdoba, Córdoba, Argentina.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fncel.2019.00122/full#supplementary-material>

## REFERENCES

- Acáz-Fonseca, E., Sanchez-Gonzalez, R., Azcoitia, I., Arevalo, M. A., and Garcia-Segura, L. M. (2014). Role of astrocytes in the neuroprotective actions of 17 $\beta$ -estradiol and selective estrogen receptor modulators. *Mol. Cell. Endocrinol.* 389, 48–57. doi: 10.1016/j.mce.2014.01.009
- Adasme, T., Haeger, P., Paula-Lima, A. C., Espinoza, I., Casas-Alarcon, M. M., Carrasco, M. A., et al. (2011). Involvement of ryanodine receptors in neurotrophin-induced hippocampal synaptic plasticity and spatial memory formation. *Proc. Natl. Acad. Sci. U S A* 108, 3029–3034. doi: 10.1073/pnas.1013580108
- Arevalo, M. A., Azcoitia, I., and Garcia-Segura, L. M. (2015). The neuroprotective actions of oestradiol and oestrogen receptors. *Nat. Rev. Neurosci.* 16, 17–29. doi: 10.1038/nrn3856
- Arevalo, M. A., Ruiz-Palmero, I., Scerbo, M. J., Acáz-Fonseca, E., Cambiasso, M. J., and Garcia-Segura, L. M. (2012). Molecular mechanisms involved in the regulation of neuritogenesis by estradiol: recent advances. *J. Steroid Biochem. Mol. Biol.* 131, 52–56. doi: 10.1016/j.jsbmb.2011.09.004
- Berridge, M. J., Bootman, M. D., and Roderick, H. L. (2003). Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* 4, 517–529. doi: 10.1038/nrm1155
- Berthois, Y., Katzenellenbogen, J. A., and Katzenellenbogen, B. S. (1986). Phenol red in tissue culture media is a weak estrogen: implications concerning the study of estrogen-responsive cells in culture. *Proc. Natl. Acad. Sci. U S A* 83, 2496–2500. doi: 10.1073/pnas.83.8.2496
- Beyer, C., and Karolczak, M. (2000). Estrogenic stimulation of neurite growth in midbrain dopaminergic neurons depends on cAMP/protein kinase A signalling. *J. Neurosci. Res.* 59, 107–116. doi: 10.1002/(sici)1097-4547(20000101)59:1<107::aid-jnr13>3.0.co;2-w
- Beyer, C., and Raab, H. (1998). Nongenomic effects of oestrogen: embryonic mouse midbrain neurones respond with a rapid release of calcium from intracellular stores. *Eur. J. Neurosci.* 10, 255–262. doi: 10.1046/j.1460-9568.1998.00045.x
- Bezprozvanny, I., Watras, J., and Ehrlich, B. E. (1991). Bell-shaped calcium-response curves of Ins(1,4,5)P<sub>3</sub>- and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature* 351, 751–754. doi: 10.1038/351751a0
- Blanco, G., Diaz, H., Carrer, H. F., and Beaugé, L. (1990). Differentiation of rat hippocampal neurons induced by estrogen *in vitro*: effects on neuritogenesis and Na, K-ATPase activity. *J. Neurosci. Res.* 27, 47–54. doi: 10.1002/jnr.490270108
- Boyan, B. D., Sylvia, V. L., Frambach, T., Lohmann, C. H., Dietl, J., Dean, D. D., et al. (2003). Estrogen-dependent rapid activation of protein kinase C in estrogen receptor-positive MCF-7 breast cancer cells and estrogen receptor-negative HCC38 cells is membrane-mediated and inhibited by tamoxifen. *Endocrinology* 144, 1812–1824. doi: 10.1210/en.2002-221018
- Brandt, D. T., Goerke, A., Heuer, M., Gimona, M., Leitges, M., Kremmer, E., et al. (2003). Protein kinase C delta induces Src kinase activity via activation of the protein tyrosine phosphatase PTP  $\alpha$ . *J. Biol. Chem.* 278, 34073–34078. doi: 10.1074/jbc.M211650200
- Brito, V. I., Carrer, H. F., and Cambiasso, M. J. (2004). Inhibition of tyrosine kinase receptor type B synthesis blocks axogenic effect of estradiol on rat hypothalamic neurones *in vitro*. *Eur. J. Neurosci.* 20, 331–337. doi: 10.1111/j.1460-9568.2004.03485.x
- Bulayeva, N. N., Wozniak, A. L., Lash, L. L., and Watson, C. S. (2005). Mechanisms of membrane estrogen receptor- $\alpha$ -mediated rapid stimulation of Ca<sup>2+</sup> levels and prolactin release in a pituitary cell line. *Am. J. Physiol. Endocrinol. Metab.* 288, E388–E397. doi: 10.1152/ajpendo.00349.2004
- Calin-Jageman, I., and Lee, A. (2008). Ca<sub>v</sub>1 L-type Ca<sup>2+</sup> channel signaling complexes in neurons. *J. Neurochem.* 105, 573–583. doi: 10.1111/j.1471-4159.2008.05286.x
- Cambiasso, M. J., and Carrer, H. F. (2001). Nongenomic mechanism mediates estradiol stimulation of axon growth in male rat hypothalamic neurons *in vitro*. *J. Neurosci. Res.* 66, 475–481. doi: 10.1002/jnr.1238
- Cambiasso, M. J., Colombo, J. A., and Carrer, H. F. (2000). Differential effect of oestradiol and astroglia-conditioned media on the growth of hypothalamic neurons from male and female rat brains. *Eur. J. Neurosci.* 12, 2291–2298. doi: 10.1046/j.1460-9568.2000.00120.x
- Cambiasso, M. J., Diaz, H., Cáceres, A., and Carrer, H. F. (1995). Neuritogenic effect of estradiol on rat ventromedial hypothalamic neurons co-cultured with homotopic or heterotopic glia. *J. Neurosci. Res.* 42, 700–709. doi: 10.1002/jnr.490420513
- Carroll, J. C., and Pike, C. J. (2008). Selective estrogen receptor modulators differentially regulate Alzheimer-like changes in female 3xTg-AD mice. *Endocrinology* 149, 2607–2611. doi: 10.1210/en.2007-1346
- Céspedes Rubio, Á. E., Pérez-Alvarez, M. J., Lapuente Chala, C., and Wandosell, F. (2018). Sex steroid hormones as neuroprotective elements in ischemia models. *J. Endocrinol.* 237, R65–R81. doi: 10.1530/joe-18-0129
- Chaban, V. V., Lakhter, A. J., and Micevych, P. (2004). A membrane estrogen receptor mediates intracellular calcium release in astrocytes. *Endocrinology* 145, 3788–3795. doi: 10.1210/en.2004-0149
- Cordey, M., Gundimeda, U., Gopalakrishna, R., and Pike, C. J. (2003). Estrogen activates protein kinase C in neurons: role in neuroprotection. *J. Neurochem.* 84, 1340–1348. doi: 10.1046/j.1471-4159.2003.01631.x
- Cullen, P. J., and Lockyer, P. J. (2002). Integration of calcium and Ras signalling. *Nat. Rev. Mol. Cell Biol.* 3, 339–348. doi: 10.1038/nrm808

- Díaz, H., Lorenzo, A., Carrer, H. F., and Cáceres, A. (1992). Time lapse study of neurite growth in hypothalamic dissociated neurons in culture: sex differences and estrogen effects. *J. Neurosci. Res.* 33, 266–281. doi: 10.1002/jnr.490330210
- Dolmetsch, R. E., Xu, K., and Lewis, R. S. (1998). Calcium oscillations increase the efficiency and specificity of gene expression. *Nature* 392, 933–936. doi: 10.1038/31960
- Dotti, C. G., Sullivan, C. A., and Banker, G. A. (1988). The establishment of polarity by hippocampal neurons in culture. *J. Neurosci.* 8, 1454–1468. doi: 10.1523/JNEUROSCI.08-04.01454.1988
- Dye, R. V., Miller, K. J., Singer, E. J., and Levine, A. J. (2012). Hormone replacement therapy and risk for neurodegenerative diseases. *Int. J. Alzheimers Dis.* 2012:258454. doi: 10.1155/2012/258454
- Falcke, M., Li, Y., Lechleiter, J. D., and Camacho, P. (2003). Modeling the dependence of the period of intracellular  $\text{Ca}^{2+}$  waves on SERCA expression. *Biophys. J.* 85, 1474–1481. doi: 10.1016/s0006-3495(03)74580-9
- Farkas, I., Sárvári, M., Aller, M., Okada, N., Okada, H., Likó, I., et al. (2012). Estrogen receptor  $\alpha$  and  $\beta$  differentially mediate C5aR agonist evoked  $\text{Ca}^{2+}$ -influx in neurons through L-type voltage-gated  $\text{Ca}^{2+}$  channels. *Neurochem. Int.* 60, 631–639. doi: 10.1016/j.neuint.2012.02.024
- Feng, Y., Wang, B., Du, F., Li, H., Wang, S., Hu, C., et al. (2013). The involvement of PI3K-mediated and L-VGCC-gated transient  $\text{Ca}^{2+}$  influx in  $17\beta$ -estradiol-mediated protection of retinal cells from  $\text{H}_2\text{O}_2$ -induced apoptosis with  $\text{Ca}^{2+}$  overload. *PLoS One* 8:e77218. doi: 10.1371/journal.pone.0077218
- Ferrari, L. F., Khomula, E. V., Araldi, D., and Levine, J. D. (2016). Marked sexual dimorphism in the role of the ryanodine receptor in a model of pain chronification in the rat. *Sci. Rep.* 6:31221. doi: 10.1038/srep31221
- Fricke, O., Kow, L. M., Bogun, M., and Pfaff, D. W. (2007). Estrogen evokes a rapid effect on intracellular calcium in neurons characterized by calcium oscillations in the arcuate nucleus. *Endocrine* 31, 279–288. doi: 10.1007/s12020-007-0034-7
- García-Segura, L. M., Arévalo, M. A., and Azcoitia, I. (2010). Interactions of estradiol and insulin-like growth factor-I signalling in the nervous system: new advances. *Prog. Brain Res.* 181, 251–272. doi: 10.1016/s0079-6123(08)81014-x
- Giatti, S., García-Segura, L. M., Barreto, G. E., and Melcangi, R. C. (2018). Neuroactive steroids, neurosteroidogenesis and sex. *Prog. Neurobiol.* doi: 10.1016/j.pneurobio.2018.06.007 [Epub ahead of print].
- Gorosito, S. V., and Cambiasso, M. J. (2008). Axogenic effect of estrogen in male rat hypothalamic neurons involves  $\text{Ca}^{2+}$ , protein kinase C, and extracellular signal-regulated kinase signaling. *J. Neurosci. Res.* 86, 145–157. doi: 10.1002/jnr.21466
- Gorosito, S. V., Lorenzo, A. G., and Cambiasso, M. J. (2008). Estrogen receptor  $\alpha$  is expressed on the cell-surface of embryonic hypothalamic neurons. *Neuroscience* 154, 1173–1177. doi: 10.1016/j.neuroscience.2008.05.001
- Hidalgo, C. (2005). Cross talk between  $\text{Ca}^{2+}$  and redox signalling cascades in muscle and neurons through the combined activation of ryanodine receptors/ $\text{Ca}^{2+}$  release channels. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 360, 2237–2246. doi: 10.1098/rstb.2005.1759
- Hidalgo, C., Donoso, P., and Carrasco, M. A. (2005). The ryanodine receptors  $\text{Ca}^{2+}$  release channels: cellular redox sensors? *IUBMB Life* 57, 315–322. doi: 10.1080/15216540500092328
- Hristov, K. L., Parajuli, S. P., Provence, A., Rovner, E. S., and Petkov, G. V. (2017). Nongenomic modulation of the large conductance voltage- and  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels by estrogen: a novel regulatory mechanism in human detrusor smooth muscle. *Physiol. Rep.* 5:e13351. doi: 10.14814/phy2.13351
- Kelly, M. J., and Levin, E. R. (2001). Rapid actions of plasma membrane estrogen receptors. *Trends Endocrinol. Metab.* 12, 152–156. doi: 10.1016/s1043-2760(01)00377-0
- Khan, M. M., Wakade, C., de Sevilla, L., and Brann, D. W. (2015). Selective estrogen receptor modulators (SERMs) enhance neurogenesis and spine density following focal cerebral ischemia. *J. Steroid Biochem. Mol. Biol.* 146, 38–47. doi: 10.1016/j.jsbmb.2014.05.001
- Khomula, E. V., Ferrari, L. F., Araldi, D., and Levine, J. D. (2017). Sexual dimorphism in a reciprocal interaction of ryanodine and IP3 receptors in the induction of hyperalgesic priming. *J. Neurosci.* 37, 2032–2044. doi: 10.1523/JNEUROSCI.2911-16.2017
- Lai, Y. J., Yu, D., Zhang, J. H., and Chen, G. J. (2017). Cooperation of genomic and rapid nongenomic actions of estrogens in synaptic plasticity. *Mol. Neurobiol.* 54, 4113–4126. doi: 10.1007/s12035-016-9979-y
- Lanner, J. T., Georgiou, D. K., Joshi, A. D., and Hamilton, S. L. (2010). Ryanodine receptors: structure, expression, molecular details, and function in calcium release. *Cold Spring Harb. Perspect. Biol.* 2:a003996. doi: 10.1101/cshperspect.a003996
- Le Mellay, V., Grosse, B., and Lieberherr, M. (1997). Phospholipase C  $\beta$  and membrane action of calcitriol and estradiol. *J. Biol. Chem.* 272, 11902–11907. doi: 10.1074/jbc.272.18.11902
- Marino, M., Pallottini, V., and Trentalancia, A. (1998). Estrogens cause rapid activation of IP3-PKC- $\alpha$  signal transduction pathway in HEPG2 cells. *Biochem. Biophys. Res. Commun.* 245, 254–258. doi: 10.1006/bbrc.1998.8413
- McCarthy, M. M. (2008). Estradiol and the developing brain. *Physiol. Rev.* 88, 91–124. doi: 10.1152/physrev.00010.2007
- McGregor, C., Riordan, A., and Thornton, J. (2017). Estrogens and the cognitive symptoms of schizophrenia: possible neuroprotective mechanisms. *Front. Neuroendocrinol.* 47, 19–33. doi: 10.1016/j.yfrne.2017.06.003
- Muchekehu, R. W., and Harvey, B. J. (2008).  $17\beta$ -estradiol rapidly mobilizes intracellular calcium from ryanodine-receptor-gated stores via a PKC-PKA-Erk-dependent pathway in the human eccrine sweat gland cell line NCL-SG3. *Cell Calcium* 44, 276–288. doi: 10.1016/j.cecc.2007.12.001
- Murphy, D. D., and Segal, M. (1996). Regulation of dendritic spine density in cultured rat hippocampal neurons by steroid hormones. *J. Neurosci.* 16, 4059–4068. doi: 10.1523/JNEUROSCI.16-13-04059.1996
- Nishizuka, M., and Arai, Y. (1981). Organizational action of estrogen on synaptic pattern in the amygdala: implications for sexual differentiation of the brain. *Brain Res.* 213, 422–426. doi: 10.1016/0006-8993(81)90247-x
- Picotto, G., Vazquez, G., and Boland, R. (1999).  $17\beta$ -oestradiol increases intracellular  $\text{Ca}^{2+}$  concentration in rat enterocytes. Potential role of phospholipase C-dependent store-operated  $\text{Ca}^{2+}$  influx. *Biochem. J.* 339, 71–77. doi: 10.1042/bj3390071
- Qiu, J., Bosch, M. A., Tobias, S. C., Grandy, D. K., Scanlan, T. S., Ronnekleiv, O. K., et al. (2003). Rapid signaling of estrogen in hypothalamic neurons involves a novel G-protein-coupled estrogen receptor that activates protein kinase C. *J. Neurosci.* 23, 9529–9540. doi: 10.1523/JNEUROSCI.23-29-09529.2003
- Quignard, J. F., Mironneau, J., Carricaburu, V., Fournier, B., Babich, A., Nurnberg, B., et al. (2001). Phosphoinositide 3-kinase  $\gamma$  mediates angiotensin II-induced stimulation of L-type calcium channels in vascular myocytes. *J. Biol. Chem.* 276, 32545–32551. doi: 10.1074/jbc.M102582200
- Reisert, I., Han, V., Lieth, E., Toran-Allerand, D., Pilgrim, C., and Lauder, J. (1987). Sex steroids promote neurite growth in mesencephalic tyrosine hydroxylase immunoreactive neurons *in vitro*. *Int. J. Dev. Neurosci.* 5, 91–98. doi: 10.1016/0736-5748(87)90054-2
- Roskoski, R. Jr. (2005). Src kinase regulation by phosphorylation and dephosphorylation. *Biochem. Biophys. Res. Commun.* 331, 1–14. doi: 10.1016/j.bbrc.2005.03.012
- Sarkar, S. N., Huang, R. Q., Logan, S. M., Yi, K. D., Dillon, G. H., and Simpkins, J. W. (2008). Estrogens directly potentiate neuronal L-type  $\text{Ca}^{2+}$  channels. *Proc. Natl. Acad. Sci. U S A* 105, 15148–15153. doi: 10.1073/pnas.0802379105
- Scerbo, M. J., Freire-Regatillo, A., Cisternas, C. D., Brunotto, M., Arevalo, M. A., García-Segura, L. M., et al. (2014). Neurogenin 3 mediates sex chromosome effects on the generation of sex differences in hypothalamic neuronal development. *Front. Cell. Neurosci.* 8:188. doi: 10.3389/fncel.2014.00188
- Seo, M. D., Enomoto, M., Ishiyama, N., Stathopoulos, P. B., and Ikura, M. (2015). Structural insights into endoplasmic reticulum stored calcium regulation by inositol 1,4,5-trisphosphate and ryanodine receptors. *Biochim. Biophys. Acta* 1853, 1980–1991. doi: 10.1016/j.bbamcr.2014.11.023
- Simoncini, T., Hafezi-Moghadam, A., Brazil, D. P., Ley, K., Chin, W. W., and Liao, J. K. (2000). Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* 407, 538–541. doi: 10.1038/35035131
- Sneyd, J., Tsaneva-Atanasova, K., Reznikov, V., Bai, Y., Sanderson, M. J., and Yule, D. I. (2006). A method for determining the dependence of calcium oscillations on inositol trisphosphate oscillations. *Proc. Natl. Acad. Sci. U S A* 103, 1675–1680. doi: 10.1073/pnas.0506135103
- Sneyd, J., Tsaneva-Atanasova, K., Yule, D. I., Thompson, J. L., and Shuttleworth, T. J. (2004). Control of calcium oscillations by membrane

- fluxes. *Proc. Natl. Acad. Sci. U S A* 101, 1392–1396. doi: 10.1073/pnas.0303472101
- Spence, R. D., Wisdom, A. J., Cao, Y., Hill, H. M., Mongerson, C. R., Stapornkul, B., et al. (2013). Estrogen mediates neuroprotection and anti-inflammatory effects during EAE through ER $\alpha$  signaling on astrocytes but not through ER $\beta$  signaling on astrocytes or neurons. *Brain Behav. Evol.* 33, 10924–10933. doi: 10.1523/jneurosci.0886-13.2013
- Striessnig, J., Pinggera, A., Kaur, G., Bock, G., and Tuluc, P. (2014). L-type  $\text{Ca}^{2+}$  channels in heart and brain. *Wiley Interdiscip. Rev. Membr. Transp. Signal.* 3, 15–38. doi: 10.1002/wmts.102
- Toran-Allerand, C. D. (1976). Sex steroids and the development of the newborn mouse hypothalamus and preoptic area *in vitro*: implications for sexual differentiation. *Brain Res.* 106, 407–412. doi: 10.1016/0006-8993(76)91038-6
- Toran-Allerand, C. D. (1980). Sex steroids and the development of the newborn mouse hypothalamus and preoptic area *in vitro*. II. Morphological correlates and hormonal specificity. *Brain Res.* 189, 413–427. doi: 10.1016/0006-8993(80)90101-8
- Toran-Allerand, C. D., Hashimoto, K., Greenough, W. T., and Saltarelli, M. (1983). Sex steroids and the development of the newborn mouse hypothalamus and preoptic area *in vitro*: III. Effects of estrogen on dendritic differentiation. *Brain Res.* 283, 97–101. doi: 10.1016/0165-3806(83)90085-8
- Vega-Vela, N. E., Osorio, D., Avila-Rodriguez, M., Gonzalez, J., Garcia-Segura, L. M., Echeverria, V., et al. (2017). L-type calcium channels modulation by estradiol. *Mol. Neurobiol.* 54, 4996–5007. doi: 10.1007/s12035-016-0045-6
- Villa, A., Vegeto, E., Poletti, A., and Maggi, A. (2016). Estrogens, neuroinflammation, and neurodegeneration. *Endocr. Rev.* 37, 372–402. doi: 10.1210/er.2016-1007
- Wong, C. K., So, W. Y., Law, S. K., Leung, F. P., Yau, K. L., Yao, X., et al. (2012). Estrogen controls embryonic stem cell proliferation via store-operated calcium entry and the nuclear factor of activated T-cells (NFAT). *J. Cell. Physiol.* 227, 2519–2530. doi: 10.1002/jcp.22990
- Wright, C. L., Schwarz, J. S., Dean, S. L., and McCarthy, M. M. (2010). Cellular mechanisms of estradiol-mediated sexual differentiation of the brain. *Trends Endocrinol. Metab.* 21, 553–561. doi: 10.1016/j.tem.2010.05.004
- Wu, T. W., Chen, S., and Brinton, R. D. (2011). Membrane estrogen receptors mediate calcium signaling and MAP kinase activation in individual hippocampal neurons. *Brain Res.* 1379, 34–43. doi: 10.1016/j.brainres.2011.01.034
- Wu, T. W., Wang, J. M., Chen, S., and Brinton, R. D. (2005).  $17\beta$ -estradiol induced  $\text{Ca}^{2+}$  influx via L-type calcium channels activates the Src/ERK/cyclic-AMP response element binding protein signal pathway and BCL-2 expression in rat hippocampal neurons: a potential initiation mechanism for estrogen-induced neuroprotection. *Neuroscience* 135, 59–72. doi: 10.1016/j.neuroscience.2004.12.027
- Yan, S., Chen, Y., Dong, M., Song, W., Belcher, S. M., and Wang, H. S. (2011). Bisphenol A and  $17\beta$ -estradiol promote arrhythmia in the female heart via alteration of calcium handling. *PLoS One* 6:e25455. doi: 10.1371/journal.pone.0025455
- Zhao, L., Chen, S., Ming Wang, J., and Brinton, R. D. (2005).  $17\beta$ -estradiol induces  $\text{Ca}^{2+}$  influx, dendritic and nuclear  $\text{Ca}^{2+}$  rise and subsequent cyclic AMP response element-binding protein activation in hippocampal neurons: a potential initiation mechanism for estrogen neurotrophism. *Neuroscience* 132, 299–311. doi: 10.1016/j.neuroscience.2004.11.054
- Zhao, X., MacBride, M. M., Peterson, B. R., Pfaff, D. W., and Vasudevan, N. (2005). Calcium flux in neuroblastoma cells is a coupling mechanism between non-genomic and genomic modes of estrogens. *Neuroendocrinology* 81, 174–182. doi: 10.1159/000087000

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Cabrera Zapata, Bollo and Cambiasso. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.